Annual Environmental Report 2019



Milford

D0342-01

CONTENTS

1 EXECUTIVE SUMMARY AND INTRODUCTION TO THE 2019 AER

- 1.1 ANNUAL STATEMENT OF MEASURES
- 1.2 TREATMENT SUMMARY
- 1.3 ELV OVERVIEW
- 1.4 LICENSE SPECIFIC REPORT INCLUDED IN AER

2 TREATMENT PLANT PERFORMANCE AND IMPACT SUMMARY

- 2.1 MILFORD (DONEGAL) WWTP TREATED DISCHARGE
 - 2.1.1 INFLUENT SUMMARY MILFORD (DONEGAL) WWTP
 - 2.1.2 EFFLUENT MONITORING SUMMARY MILFORD (DONEGAL) WWTP -
 - 2.1.3 Ambient Monitoring Summary for The Treatment Plant Discharge -
 - 2.1.4 OPERATIONAL REPORTS SUMMARY FOR MILFORD (DONEGAL) WWTP
 - 2.1.5 SLUDGE/OTHER INPUTS TO MILFORD (DONEGAL) WWTP

3 COMPLAINTS AND INCIDENTS

- 3.1 COMPLAINTS SUMMARY
- 3.2 REPORTED INCIDENTS SUMMARY
 - 3.2.1 SUMMARY OF INCIDENTS
 - 3.2.2 SUMMARY OF OVERALL INCIDENTS

4 INFRASTRUCTURAL ASSESSMENT AND PROGRAMME OF IMPROVEMENTS

- 4.1 STORM WATER OVERFLOW IDENTIFICATION AND INSPECTION REPORT
 - 4.1.1 SWO IDENTIFICATION AND INSPECTION SUMMARY REPORT
- 4.2 REPORT ON PROGRESS MADE AND PROPOSALS BEING DEVELOPED TO MEET THE IMPROVEMENT PROGRAMME REQUIREMENTS
- 4.2.1 Specified Improvement Programme Summary
- 4.2.2 IMPROVEMENT PROGRAMME SUMMARY
- 4.2.3 SEWER INTEGRITY RISK ASSESSMENT

5 LICENCE SPECIFIC REPORTS

- 5.1 PRIORITY SUBSTANCES ASSESSMENT
- 5.2 SMALL STREAM RISK SCORE ASSESSMENT
- 6 CERTIFICATION AND SIGN OFF
 - 6.1 SUMMARY OF AER CONTENTS

7 APPENDIX

7.1 SMALL STREAM RISK SCORE ASSESSMENT

1 EXECUTIVE SUMMARY AND INTRODUCTION TO THE 2019 AER

This Annual Environmental Report has been prepared for D0342-01, Milford, in Donegal in accordance with the requirements of the wastewater discharge licence for the agglomeration. Specified reports where relevant are included as an appendix to the AER.

1.1 ANNUAL STATEMENT OF MEASURES

A summary of any improvements undertaken is provided where applicable. Milford WwTW is included on the Irish Water Capital Investment Programme 2018 - 2021.

1.2 TREATMENT SUMMARY

The agglomeration is served by a wastewater treatment plant(s)

• MILFORD (DONEGAL) WWTP with a Plant Capacity PE of 920, the treatment type is 2 - Secondary treatment

1.3 ELV OVERVIEW

The overall compliance of the final effluent with the Emission Limit Values (ELVs) is shown below. More detailed information on the below ELV's can be found in Section 2.

Discharge Point Reference	Treatment Plant	Discharge Type	Compliance Status	Parameters failing if relevant
TPEFF0600D0342SW001	MILFORD (DONEGAL) WWTP	Treated	Non-Compliant	Ammonia-Total (as N) mg/l BOD, 5 days with Inhibition (Carbonaceous BOD) mg/l COD-Cr mg/l ortho-Phosphate (as P) - unspecified mg/l Suspended Solids mg/l

1.4 LICENCE SPECIFIC REPORTING INCLUDED IN AER

Assessment / Report	Included in AER
Small Stream Risk Score Assessment	Yes

2 TREATMENT PLANT PERFORMANCE AND IMPACT SUMMARY

2.1 MILFORD (DONEGAL) WWTP - TREATED DISCHARGE

2.1.1 INFLUENT MONITORING SUMMARY - MILFORD (DONEGAL) WWTP

A summary of influent monitoring for the treatment plant is presented below. This monitoring is primarily undertaken in order to determine the overall efficiency of the plant in removing pollutants from the raw wastewater.

Parameters	Number of Samples	Annual Max	Annual Mean
Total Nitrogen mg/l	11	34.9	19.24
BOD, 5 days with Inhibition (Carbonaceous BOD) mg/I	11	130	67.84
Suspended Solids mg/l	11	154	72.79
COD-Cr mg/l	11	231	146.18
Total Phosphorus (as P) mg/l	11	4.28	2.05
Hydraulic Capacity	N/A	3966	1161

If other inputs in the form of sludge / leachate are added to the WWTP then these are included in Section 2.1.5 if applicable.

Significance of Results:

The annual mean hydraulic loading is greater than the peak Treatment Plant Capacity. The annual maximum hydraulic loading is greater than the peak Treatment Plant Capacity. Further details on the plant capacity and efficiency can be found under the sectional 'Operational Performance Summary'.

2.1.2 EFFLUENT MONITORING SUMMARY - TPEFF0600D0342SW001

Parameter	WWDL ELV (Schedule A)	ELV with Condition 2 Interpretation included Note 1	Interim % reduction from influent concentration	Number of sample results	Number of exceedances	Number of with Condition 2 Interpretation included	Annual Mean	Overall Compliance (Pass/Fail)
COD-Cr mg/l	125	250	N/A	11	5	1	114.86	Fail
Suspended Solids mg/l	25	62.5	N/A	11	6	3	52.38	Fail
BOD, 5 days with Inhibition (Carbonaceous BOD) mg/l	10	20	N/A	11	8	6	28.17	Fail
pH pH units	9	9	N/A	11	N/A	N/A	7.23	Pass
Ammonia-Total (as N) mg/l	0.65	1.3	N/A	11	11	11	14.52	Fail
ortho-Phosphate (as P) - unspecified mg/l	0.34	0.68	N/A	11	11	8	1.3	Fail
Total Nitrogen mg/l	N/A	N/A	N/A	11	N/A	N/A	18.15	
Conductivity 20 C µS/cm	N/A	N/A	N/A	11	N/A	N/A	455.71	
Total Phosphorus (as P) mg/l	N/A	N/A	N/A	11	N/A	N/A	1.81	

Notes:

1 - This represents the Emission Limit Values after the Interpretation provided for under Condition 2 of the licence is applied

Cause of Exceedance(s):

Plant is overloaded

Significance of Results:

The WWTP is non-compliant with ELV's set in the Wastewater Discharge Licence. The impact on the receiving waters is assessed further in Section 2.

2.1.3 AMBIENT MONITORING SUMMARY FOR THE TREATMENT PLANT DISCHARGE TPEFF0600D0342SW001

A summary of monitoring from ambient monitoring points associated with the wastewater discharge is provided in the sections below. For discharges to rivers upstream (U/S) and downstream (D/S) location data is provided. For other ambient points in lakes, coastal or transitional waters, monitoring data from the most appropriate monitoring station is selected.

The table below provides details of ambient monitoring locations and details of any designations as sensitive areas.

Ambient Monitoring Point from WWDL (or as agreed with EPA)	Irish Grid Reference	River Station Code	Bathing Water	Drinking Water	FWPM	Shellfish	WFD Status
Upstream	219277, 426509	RS39M010150	No	No	No	No	Poor
Downstream	218606, 424918	RS39M010300	No	No	No	No	Poor

The table below provides a summary of monitoring results for designated ambient monitoring points. The upstream and downstream annual mean values are shown (mg/l), and the difference between both monitoring stations is given as a percentage of the Environmental Quality Standard (EQS) where relevant.

Parameter Name	Upstream Monitoring Point Location	Upstream Monitoring Point Annual Mean	Downstream Monitoring Point Location	Downstream Monitoring Point Annual Mean	EQS	% of EQS
BOD - 5 days (Total) mg/l	RS39M010150	1.944	RS39M010300	3.338	1.5	92.9
Ammonia-Total (as N) mg/l	RS39M010150	0.072	RS39M010300	0.539	0.065	719.3
ortho-Phosphate (as P) - unspecified mg/l	RS39M010150	0.028	RS39M010300	0.102	0.035	214.1
Total Phosphorus (as P) mg/l	RS39M010150	0.042	RS39M010300	0.189		
Orthophosphate (MRP) filtered (As P) mg/I	RS39M010150	0.025	RS39M010300	0.09		
Dissolved Oxygen % Saturation	RS39M010150	97.078	RS39M010300	67.443		
Conductivity 20 C µS/cm	RS39M010150	272.556	RS39M010300	268.556		
Total Nitrogen mg/l	RS39M010150	1.683	RS39M010300	2.62		
Temperature °C	RS39M010150	11.888	RS39M010300	10.938		
pH pH units	RS39M010150	7.411	RS39M010300	6.986		
Suspended Solids mg/l	RS39M010150	8.25	RS39M010300	15		

Significance of Results:

The WWTP discharge was not compliant with the ELV's set in the wastewater discharge licence.

The ambient monitoring results does not meet the required EQS. The EQS relates to the Oxygenation and Nutrient Conditions set out in the Surface Water Regulations 2009.

Based on ambient monitoring results a deterioration in Ammonia, Ortho-Phosphate, BOD and Total Phosphorus, concentrations downstream of the effluent discharge is noted.

A deterioration in water quality has been identified, however it is not known if it or is not caused by the WWTP.

Other causes of deterioration in water quality in the area are unknown.

The discharge from the wastewater treatment plant does not have an observable negative impact on the Water Framework Directive status.

2.1.4 OPERATIONAL PERFORMANCE SUMMARY - MILFORD (DONEGAL) WWTP

2.1.4.1 Treatment Efficiency Report - MILFORD (DONEGAL) WWTP

Treatment efficiency is based on the removal of key pollutants from the influent wastewater by the treatment plant. In essence the calculation is based on the balance of load coming into the plant versus the load leaving the plant. The efficiency is presented as a percentage removal rate.

A summary presentation of the efficiency of the treatment process including information for all the parameters specified in the licence is included below:

Parameter	Influent mass loading (kg/year)	Effluent mass emission (kg/year)	Efficiency (% reduction of influent load)
TN	488	460	5.67
ТР	52	46	12
SS	1847	1329	28
cBOD	1721	715	58
COD	3708	2914	21

Note: The above data is based on sample results for the number of dates reported

2.1.4.2 Treatment Capacity Report Summary - MILFORD (DONEGAL) WWTP

Treatment capacity is an assessment of the hydraulic (flow) and organic (the amount of pollutants) load a treatment plant is designed to treat versus the current loading of that plant.

MILFORD (DONEGAL) WWTP	
Peak Hydraulic Capacity (m³/day) - As Constructed	621
DWF to the Treatment Plant (m ³ /day)	207
Current Hydraulic Loading - annual max (m³/day)	3966
Average Hydraulic loading to the Treatment Plant (m³/day)	1161
Organic Capacity (PE) - As Constructed	920
Organic Capacity (PE) - Collected Load (peak week) ^{Note1}	1678
Organic Capacity (PE) - Remaining	0
Will the capacity be exceeded in the next three years? (Yes/No)	Yes

Nominal design capacities can be based on conservative design principles. In some cases assessment of existing plants has shown organic capacities significantly higher than the nominal design capacity. Accordingly plants that appear to be overloaded when comparing a collected peak load with the nominal design capacity can be fully compliant due to the safety factors in the original design.

2.1.5 SLUDGE / OTHER INPUTS - MILFORD (DONEGAL) WWTP

'Other inputs' to the waste water treatment plant are summarised in table below

Input type	Quantity	Unit	P.E.	% of load to WWTP	Included in Influent Monitoring (Y/N)?	Is there a leachate/sludge acceptance procedure for the WWTP?	Is there a dedicated leachate/sludge acceptance facility for the WWTP? (Y/N)		
There is	There is no Sludge and Other Input data for the Treatment Plant included in the AER.								

3 COMPLAINTS AND INCIDENTS

3.1 COMPLAINTS SUMMARY

A summary of complaints of an environmental nature is included below.

Number of Complaints	Nature of Complaint	Number Open Complaints	Number Closed Complaints
There were no relevant environme	ental complaints in 2019.		

3.2 REPORTED INCIDENTS SUMMARY

Environmental incidents that arise in an agglomeration are reported on an on-going basis in accordance with our waste water discharge licences. Where an incident occurs and it is reportable under the licence, it is reported to the Environmental Protection Agency through their Environmental Data Exchange Network, or in some instances by telephone. Some incidents which arise in the agglomeration are recorded by Irish Water but may not be reportable under our licence for example where the incident does not have an impact on environmental performance.

A summary of reported incidents is included below.

3.2.1 SUMMARY OF INCIDENTS

Incident Type	Cause	No. of incident occurrences	Recurring (Y/N)	Closed (Y/N)
Breach of ELV	WWTP operating above capacity	1	Yes	No

3.2.2 SUMMARY OF OVERALL INCIDENTS

Question	Answer
Number of Incidents in 2019	1
Number of Incidents reported to the EPA via EDEN in 2019	1
Explanation of any discrepancies between the two numbers above	N/A

4 INFRASTRUCTURAL ASSESSMENTS AND PROGRAMME OF IMPROVEMENTS

4.1 STORM WATER OVERFLOW IDENTIFICATION AND INSPECTION REPORT

A summary of the operation of the storm water overflows and their significance where known is included below:

4.1.1 SWO IDENTIFICATION

WWDL Name / Code for Storm Water Overflow	Irish Grid Ref.	Included in Schedule A4 of the WWDL	Significance of the overflow(High / Medium / Low)	Assessed against DoEHLG Criteria	No. of times activated in 2019 (No. of events)	Total volume discharged in 2019 (m3)	Monitoring Status
SW3	219287, 426533	Yes	Unknown	Not Meeting	Unknown	Unknown	Not Monitored
твс	твс	No	Unknown	Not yet Assessed	Unknown	Unknown	Not Monitored
SW2	219237, 426535	Yes	Low	Meeting	Unknown	Unknown	Not Monitored

SWO Summary	
How much sewage was discharged via SWOs in the agglomeration in the year (m3)?	Unknown
Is each SWO identified as not meeting DoEHLG Guidance included in the Programme of Improvements?	Yes
The SWO Assessment included the requirements of relevant of WWDL schedules?	Yes

SWO Summary

Have the EPA been advised of any additional SWOs / changes to Schedule C3 and A4 under Condition 1.7?

4.2 REPORT ON PROGRESS MADE AND PROPOSALS BEING DEVELOPED TO MEET THE IMPROVEMENT PROGRAMME REQUIREMENTS.

4.2.1 SPECIFIED IMPROVEMENT PROGRAMME SUMMARY

A wastewater discharge licence may require a number of reports on specific subject areas to be prepared for the agglomeration in question. These reports are submitted to the EPA as part of the Annual Environmental Report. This section provides list of the various reports required for this agglomeration and a brief summary of their recommendations.

Specified Improvement Programmes (under Schedule A and C of WWDL)	Description	Licence Schedule	Licence Completion Date	Date Expired? (N/NA/Y)	Status of Works	Timeframe for Completing the Work	Comments
D0342-SIP:01	Infiltration programme - diversion of storm water from the sewer collection network	С	31/12/2012	Yes	Not Started		The improvement programme will be reviewed by Irish Water to assess the works required to comply with the licence condition on a prioritised basis
D0342-SIP:02	Installation of storm water storage tank	С	31/12/2017	Yes	At Planning Stage	22/09/2023	

Specified Improvement Programmes (under Schedule A and C of WWDL)	Description	Licence Schedule	Licence Completion Date	Date Expired? (N/NA/Y)	Status of Works	Timeframe for Completing the Work	Comments
D0342-SIP:03	Redesign WWTP inlet works for better flow control	С	31/12/2012	Yes	Works Completed		
D0342-SIP:05	WWTP expansion and upgrade to provide tertiary treatment	С	31/12/2017	Yes	At Planning Stage	22/09/2023	Ramelton, Milford & Rathmullan will be served by 1 No. Secondary WWTP and have one discharge point.
D0342-SIP:04	Upgrading of Storm Water Overflows to comply with the criteria outlined in the DoECLG 'Procedures and Criteria in relation to Storm Water overflows, 1995'	С	31/12/2017	Yes	At Planning Stage	31/12/2023	

A summary of the status of any improvements identified by under Condition 5.2 is included below.

4.2.2 IMPROVEMENT PROGRAMME SUMMARY

Improvement	Improvement Description / or any Operational	Improvement	Expected Completion	Comments
Identifier	Improvements	Source	Date	
There are no Improven	nents Programme for this Agglomeration.			

4.2.3 SEWER INTEGRITY RISK ASSESSMENT

The utilisation of multiple capital maintenance programmes and the outputs of the workshops with the Local Authority Operations Staff held under the programme can be used to satisfy the requirements of Condition 5 regarding network integrity. Improvement works identified by way of these programmes and workshops will be included in the Improvements Summary Table.

5 LICENCE SPECIFIC REPORTS

A wastewater discharge licence may require a number of reports on specific subject areas to be prepared for the agglomeration in question. These reports are submitted to the EPA as part of the Annual Environmental Report. This section provides list of the various reports required for this agglomeration and a brief summary of their recommendations.

5.a Licence Specific Reports Summary Table

Licence Specific Report	Required by licence	Year included in AER	Included in this AER	Reference to relevant section of AER
Priority Substances Assessment	Yes	2015	No	
Small Stream Risk Score Assessment	Yes	2016	Yes	5.2

5.1 PRIORITY SUBSTANCES ASSESSMENT

The Priority Substances Assessment Report has been included in the AER 2015

5.2 SMALL STREAM RISK SCORE ASSESSMENT

The Small Stream Risk Score Assessment Report is included in Appendix 7.1 - Small Stream Risk Score Assessment. A summary of the findings of this report is included below.

Parameter	Value
Condition 5 Improvement Programme Reference	Ref. 4.2 Specified Improvement Programme

Parameter	Value
Does SSRS indicate discharges are posing a pollution risk?	Yes
Does improvement programme include any procedural and/or infrastructal works?	Yes
Downstream SSRS Water Quality Risk	N/A
SSRS Required?	Yes
Upstream SSRS Water Quality Risk	N/A
What is Downstream SSRS?	Q2-3
What is Upstream SSRS?	Q3

6 CERTIFICATION AND SIGN OFF

6.1 SUMMARY OF AER CONTENTS

Parameter	Answer
Does the AER include an Executive Summary?	Yes
Does the AER include an assessment of the performance of the Waste Water Works (i.e. have the results of assessments been interpreted against WWDL requirements and or Environmental Quality Standards)?	Yes
Is there a need to advise the EPA for consideration of a Technical Amendment / Review of the licence?	Yes
List reason e.g. additional SWO identified	Ramelton, Milford & Rathmullan will be served by 1 No. Secondary WWTP and have one discharge point. A Licence review will be submitted to accommodate the combination of the various agglomerations.
Is there a need to request/advise the EPA of any modification to the existing WWDL with respect to condition 4 changes to monitoring location, frequency etc	No
List reason e.g. changes to monitoring requirements	N/A
Have these processes commenced?	No
Are all outstanding reports and assessments from previous AERs included as an appendix to this AER	No

I certify that the information given in this Annual Environmental Report is truthful, accurate and complete:

Signed: Date: 16/06/2020

This AER has been produced by Irish Water's Environmental Information System (EIMS) and has been electronically signed off in that system for and on behalf of ,

Katherine Walshe

Acting Head of Environmental Regulation.

7 APPENDIX

Appendix

Appendix 7.1 - Small Stream Risk Score Assessment


Station no.	maggie's Burn Milford		Grid (6 figure):
Station no. 19250	1388	Location: UpStream Stream Order:	Stream flow:
Field Ch	emistry	Modifications: Y(n)Canalised-wirdened bank ero	Ision Riffle/Glide
0% Ι\ριη ΟΟ	95.6	arterial drainage Dominant Types:	Slow flow
Temp (°C)	10.4	Bedrock	
Conductivity	271	Boulder (>128mm) Cobble (32-128mm)	
H	736	Gravel (8-32mm)	
Bank width (cm)	350	Fine Gravel (2-8mm)	
Vet width (cm)	250	- Sand (0.25-2mm) Silt (<0.25mm)	
vg Depth (cm)	70	Slope: Low (Hedium) High Very High	Arrest of the second
Staff gauge	10		Shading: High Moderate Low None
Velocity	Colour	Geology: Calcareous Siliceou -M x+d	
Torrential	None	Substratum Condition: Calcareous Compacted	Cattle access Y: upstream downstream o
Moderate	(Slight)	Loose - Normal Substratum:	
Slow	High	Stoney bottom Muddy bottom slud over stones	Photo: Y / N
Very slow		Degree of siltation: Clean Slight Moderate Hea	
Clarity	Discharge	Depth of mud: (Ione) - 1cm: 1-5cm - 5-10cm -	
Very clear	Flood	Litter: None Present Moderate Abundant	
Clear	Normal		Sources Fire Fire
Sightiy turbid	Low	Filamentous Algae: None Present Moderate Abundant	Sewage Fungus: None – Present – Moderate – Abundarit
Highly turbid	Very Low	Main land use u/s: Sample	Sampled in Minutes:
	Dry	Pasture Urhan retained	: Flond net x XIO
	Recent Flood	Bog Tillage (Y) N ' Forestry Other	Stork wash x ×10
			Weed stylep x
General Commen	its:		
The macroinvertebr	ates are divided int	Macroinvertebrate Composition	Relative Abundance
The macroinvertebr Group 1 = 1	ates are divided int Ephemeroptera (3-ti	o the following 5 specific groups: ails) – note that tails may be damaged during samp	Relative Abundance 1-5
The macroinvertebr Group 1 = 1	ates are divided int Ephemeroptera (3-ti Piecoptera (2-tails)	o the following 5 specific groups:	Relative Abundance
The macroinvertebr Group 1 = 1 Group 2 = 1 Group 3 = 1 Group 4 = 0	ates are divided int Ephemeroptera (3-ta Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda	o the following 5 specific groups: ails) – note that tails may be damaged during samp	Pling 1-5 6-20 21-50 51-100
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Group 5 = 2	ates are divided int Ephemeroptera (3-t Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sallus</i>	o the following 5 specific groups: ails) – note that tails may be damaged during samp - note that tails may be damaged during sampling	Dling 1-5 6-20 21-50 51-100 101 +
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-t Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sallus</i>	o the following 5 specific groups: ails) – note that tails may be damaged during samp - note that tails may be damaged during sampling a Oligochaeta and Diptera) asa and relative abundance of each macroinvertebra	ate group below." (Abundance - Ab)
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-t Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sallus</i>	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Diptera - Oligochaeta and Diptera - Add relative abundance of each macromivertebra - Ecdy onurus Ab Plecoptera;	bling 1-5 6-20 21-50 51-100 101 + ate group below (Abundance – Ab) Leuctra Ab
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-t Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sallus</i>	o the following 5 specific groups: ails) – note that tails may be damaged during samp - note that tails may be damaged during sampling a Oligochaeta and Diptera) asa and relative abundance of each macroinvertebra	ate group below." (Abundance - Ab)
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-t Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sallus</i>	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Diptera - Oligochaeta and Diptera - Add relative abundance of each macromvertebra - Ecdy onurus Ab - Rhithrogena Ab	bling 1-5 6-20 21-50 51-100 101 + ate group below (Abundance - Ab) <i>Leuctra</i> Ab <i>Isoper/a</i> Ab
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-t Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sallus</i>	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Diptera) axa and relative abundance of each macromivertebra - Ecdyonurus Ab - Rhithrogena Ab - Heptagenia Ab - Ephemerella Ab	ate group below = (Abundance - Ab) Relative Abundance 1-5 6-20 21-50 51-100 101 + Leuctra Ab Isoper/a Ab Protonemura Ab
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-t Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sallus</i> a total number of b	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Dipteral axa and relative abundance of each macromvertebra - Ecdy onurus Ab - Rhithrogena Ab - Heptagenia Ab - Caenis Ab	oling 1-5 6-20 21-50 51-100 101+ ate group below (Abundance = Ab) <i>Leuctra</i> Ab <i>Isoper/a</i> Ab <i>Protonemura</i> Ab <i>Amphinemura</i> Ab
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-ta Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda Asallus ne total number of ta	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling a Oligochaeta and Diptera) asa and relative abundance of each macroinvertebra <i>Ecdy onurus</i> Ab <i>Rhithrogena</i> Ab <i>Heptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab	Bling Abundance 1-5 6-20 21-50 51-100 101 + ate group below (Abundance - Ab) <i>Leuctra</i> Ab <i>Isoper/a</i> Ab <i>Protonemura</i> Ab <i>Anphinemura</i> Ab <i>Per/a</i> Ab <i>Dinocras</i> Ab
The macroinvertebr Group 1 = 6 Group 2 = 7 Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-ta Piecoptera (2-tails) Trichoptera G.OL D (Gastropoda Asallus ne total number of ta	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Dipteral axa and relative abundance of each macroinvertebra - Ecdy onurus Ab - Rhithrogena Ab - Heptagenia Ab - Caenis Ab - Caenis Ab - Dipterar danica Ab	bling Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoper/a Ab Protonentura Ab Amphinemura Ab Per/a Ab Dinocras Ab Other Piecop Ab
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 C Group 5 Calculate th	ates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sellus</i> ne total number of h <i>F</i>	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Dipteral axa and relative abundance of each macromvertebra <i>Ecdy onurus</i> Ab <i>Rhithrogena</i> Ab <i>Heptagenia</i> Ab <i>Caenis</i> Ab <i>Caenis</i> Ab <i>Caenis</i> Ab <i>Caenis</i> Ab <i>Chemeral danica</i> Ab Other Ephem Ab	Pling 1-5 6-20 21-50 51-100 101 + ate group below (Abundance – Ab) <i>Leuctra</i> Ab <i>Isoperia</i> Ab <i>Protonemura</i> Ab <i>Amphinemura</i> Ab <i>Peria</i> Ab <i>Cinocras</i> Ab Other Piecop Ab
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails)) Trichoptera G.OL.D. (Gastropoda A <i>sellus</i> total number of 1 <i>Ej</i> xa	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Diptera) - Asa and relative abundance of each macromvertebra - Ecdy onurus Ab - Rhithrogena Ab - Heptagenia Ab - Ephemerella Ab - Caenis Ab - Plecoptera: - Plecoptera:	Bling 1-5 6-20 21-50 51-100 101 + ate group below (Abundance - Ab) <i>Lauctra</i> Ab <i>Isoperia</i> Ab <i>Protonemura</i> Ab <i>Protonemura</i> Ab <i>Peria</i> Ab <i>Peria</i> Ab <i>Dinocras</i> Ab Other Piecop Ab
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Calculate th Ephemeroptera:	ates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G.OL D (Gastropoda A <i>sellus</i> ne total number of h <i>F</i>	o the following 5 specific groups: ails) - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - note that tails may be damaged during sampling - Ridthat tails may be damaged during sampling - Ecdy onurus Ab - Plecoptera: - Rhithrogena Ab - Plecoptera: - Caenis Ab - Plecoptera - Other Ephem Ab - Plecoptera - Other Ephem Ab - Plecoptera - Other Ephem Ab - Other Ephem Ab - Other Ephem Ab - Other Ephem Ab	Abundance 1-5 6-20 21-50 51-100 101 + ate group belox. (Abundance - Ab) <i>Leuctra</i> Ab <i>Isoperla</i> Ab <i>Protonemura</i> Ab <i>Perla</i> Ab <i>Perla</i> Ab <i>Dinocras</i> Ab Other Plecop Ab Other Plecop Ab
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera G.OL D (Gastropoda Asellus te total number of ta <i>E</i> E Xa C Total R Hydropsychi	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Dipteral axa and relative abundance of each macroinvertebra <i>Ecdy onurus</i> Ab <i>Ecdy onurus</i> Ab <i>Rhithrogena</i> Ab <i>Heptagenia</i> Ab <i>Caenis</i> Ab	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperia Ab Protonemura Ab Amphinemura Ab Peria Ab Dinocras Ab Other Plecop Ab Other Plecop Ab Other Plecop Ab Chironomidae (D) Ab Simulidae (D) Ab 3 Few/Low
Group 1 = F Group 2 = F Group 3 = 7 Group 4 = 0 Group 5 = 7 Calculate th	ates are divided int Ephemeroptera (3-ta Plecoptera (2-tals) Trichoptera G.OL D (Gastropod a Ase//us ne total number of b total number of b Ej xa o total R Hydropsychi Polycentropodi <i>Rhy acop</i> Philopotami	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling a Oligochaeta and Dipteral a cand relative abundance of each macroinvertebra <i>Ecdy onurus</i> Ab <i>Ecdy onurus</i> Ab <i>Rhithrogena</i> Ab <i>Heptagenia</i> Ab <i>Caenis</i> Ab	Ding Abundance 1.5 6.20 21-50 51-100 101 + ate group below (Abundance - Ab) <i>Leuctra</i> Ab <i>Isoperia</i> Ab <i>Protonemura</i> Ab <i>Protonemura</i> Ab <i>Peria</i> Ab <i>Dinocras</i> Ab Other Piecop Ab Chironomidae (D) Ab Simulidae (D) Ab Common/
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Calculate th Ephemeroptera:	ates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G.OL D (Gastropoda Ase//us total number of la total number of la Ej xa ST Total R Hydropsychir Polycentropodi <i>Rh</i>) acop Philopotami Limnephili	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Dipteral - Oligochaeta and Dipteral - Asa and relative abundance of each macromvertebra - Ecdy onurus Ab - Ecdy onurus Ab - Rhithrogena Ab - Heptagenia Ab - Ephemeralia Ab - Caenis Ab - Other Ephem Ab - elative Abundance - Other Ephem Ab - elative Abundance - Other Ephem Ab - elative Abundance - Other Ephem Ab - Potamopy rgus (G) Ab - Ab	Pling Relative Abundance 1-5 6-20 21-50 51-100 101 + 101 + ate group below (Abundance - Ab) Leuctra Ab Loger/a Ab Protonemura Ab Aniphinemura Ab Per/a Ab Dinocras Ab Other Plecop Ab Other Plecop Ab Other Plecop Ab Chironomidae (D) Ab Absent Simulidae (D) Ab Absent Simulidae (D) Ab Common, Tipulidae (D) Ab Numerous
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Piecoptera (2-tails)) Trichoptera G.OL D (Gastropoda Aseilus ne total number of b total number of b <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i>	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Diptera) aka and relative abundance of each macromvertebra Ecdy onurus Ab Rhithrogena Ab Heptagenia Ab Caenis Ab Caenis Ab Other Ephem Ab elative Abundance dae Ab G.OL.D: Ly minaea (G) Ab planorbis (G) Ab Ab Caenis (G) Ab Ab Caenis (G) Ab Ab Caenis (G) Ab Ab Caenis (G) Ab Ab Caenis (G) Ab Caenis (C) Ab	Dling Abundance 1-5 6-20 21-50 51-100 101+ ate group below (Abundance – Ab) <i>Lauctra</i> Ab <i>Isoperla</i> Ab <i>Protonemura</i> Ab <i>Protonemura</i> Ab <i>Perla</i> Ab <i>Dinocras</i> Ab Other Plecop Ab Other Plecop Ab Other Plecop Ab Other Plecop Ab Other Plecop Ab Chironomidae (D) Ab <i>Simuliidae</i> (D) Ab <i>Simuliidae</i> (D) Ab <i>Simuliidae</i> (D) Ab <i>Simuliidae</i> (D) Ab <i>Ceratopogonidae</i> (D) Ab
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G.OL D (Gastropoda Asellus total number of la total number of la Ej xa C Total R Hydropsychi Polycentropodi <i>Rhy acop</i> Philopotami Limnephili Sericostomati Giossosomati	o the following 5 specific groups: ails) – note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Diptera) axa and relative abundance of each macromvertebra Ecdy onurus Ab Rhithrogena Ab Heptagenia Ab Caenis Ab Caenis Ab Plecoptera: Rhithrogena Ab Caenis Ab Caenis Ab Plecoptera: Total no. of Taxa dae Ab G.OL.D: Lynnaea (G) Ab Planorbis (G) Ab Ancylus (G) Ab Caenis Ab	Ding Abundance 1.5 6.20 21.50 51.100 101.4 ate group below (Abundance - Ab) <i>Leuctra</i> Ab <i>Isoperla</i> Ab <i>Protonemura</i> Ab <i>Amph.nemura</i> Ab <i>Perla</i> Ab <i>Dinocras</i> Ab Other Piecop Ab Other Piecop Ab Other Piecop Ab Chironomidae (D) Ab <i>Simulidae</i> (D) Ab
The macroinvertebr Group 1 = E Group 2 = F Group 3 = C Group 4 = C Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Piecoptera (2-tails)) Trichoptera G.OL D (Gastropoda Aseilus ne total number of b total number of b <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i> <i>t</i>	o the following 5 specific groups: ails) - note that tails may be damaged during sampling - note that tails may be damaged during sampling - Oligochaeta and Diptera) axa and relative abundance of each macromvertebra Ecdy onurus Ab Rhithrogena Ab Heptagenia Ab Ephemerella Ab Caenis Ab Plecoptera: Noter Ephem Ab elative Abundance dae Ab G.OL.D: Lymnaea (G) Ab axa Ab Pianorbis (G) Ab dae Ab dae Ab dae Ab dae Ab Ab Caenis Ab Pianorbis (G) Ab Caenis (C) Ab Caen	Abundance 1.5 6.20 21-50 51-100 101 + ate group below (Abundance - Ab) <i>Leuctra</i> Ab <i>Isoperia</i> Ab <i>Protonemura</i> Ab <i>Protonemura</i> Ab <i>Protonemura</i> Ab <i>Peria</i> Ab <i>Dinocras</i> Ab Other Plecop Ab Other Plecop Ab Other Plecop Ab Chironomidae (D) Ab <i>Simulidae</i> (D) Ab

NOTE *Baetis* is an Ephemeropteran and is the most commonly occurring invertebrate genus in streams in Ireland. It is vital that *Baetis* is not counted in SSRS. See Appendix B for more details on how to identify *Baetis*.

100

Г 50


의 111 位 **Step 1.** Calculate the Index Score by circling the appropriate box representing the total number of taxa and the total abundance calculated from *each macroinvertebrate group* calculated from page 1 of the recording sheet and enter in to the boxes in Step 2.



Step 3. Calculate the Total Index Score, the Average Index Score and the SSR Score using the boxes below



Step 4. Assess the stream by comparing the final SSR score with the categories below and tick the appropriate box




River: D/S Maggie's Burn Station no. 19250@ 1389		Location: Do	whetream	Grid (6 figure): Stream flow:	
			analised oldened bank erosion	Piffle	
Field Ch	emistry 8217 %	arterial drainage	Lanaised videned bank erosion	Riffle/Glide	
00 mg/l	021110	Dominant Types:		Pour nom	
emp (°C)	11°C	Bedrock.			
Conductivity		Boulder (>128mm) Cobble (32-128mm)			
H	305	Gravel (8-32mm)		And a second	
	-	Fine Gravel (2-8nim)			
Bank width (cm)	180	Sand (0.25-2mm)			
Vet width (cm)	120	(ilt)-0.25mm)			
Vg Depth (cm)	80	Slope: Low Medium	n - High - Very High	Shading: High - Móderati	lo : Nona
itaff gauge Velocity	Colour	Geology: Calcareous	Siliceou -Mix-d	Shaung: Byn - Mouerau	U.V. = 110116
Torrential	None	Substratum Condit	ion: Calcareous-Compacted	Cattle access Y upstream	downstream c
Fast	Sligto	Loose - Normal			
Hoderau	Moderate	Substratum:		~	
Stavy	High		y bottoe Hud over stones	Photo: Y (N)	
Very slow	Discharge		: Clean Slight-Moderate Heavy		
Clarity Very clear	Flood	Depth of mud: Non	e <1cm (1-5cm) 5-10cm - 10cm		
Clear	Normal	Litter: Noné - Presa	nt (Inderate Abundant		
Clean	a annai	1		Sewage Fongus:	
Slightly turb d	Lon	Filamentous Algae None = Present = Mo		None - (resent) - Moderate	Abundant
Highly turbid	Very Low	Main Land use u/s:	Sample	Sampled in Minutes:	
2	Dry	Pasture	Urban retained:	Pond net x	
	Recent Flood	Bog	Tillage (Y) N	Stone wash ×	
		Forestry	Other		
General Commer	nts:	Macroinvertebra	ate Composition	Weed sweep ×	Relative
Group 1 🕾	rates are divided int Ephemeroptera (3-t	o the following 5 specificants) = note that tails ma	iy be damaged during sampling	Weed sweep x	Relative Abundance
The macroinverteb Group 1 = Group 2 =	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails)	o the following 5 specificants) = note that tails ma	c groups:	Weed sweep x	Abundance 1-5 6-20
The macroinverteb Group 1 = Group 2 = Group 3	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera	o the following 5 specifi ails) – note that tails ma - note that tails may be	c groups: iy be damaged during sampling damaged during sampling	Weed sweep x	Abundance 1-5 6-20 21-50
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 =	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL-D (Gastropode <i>Asellus</i>	o the following 5 specifi ails) – note that tails ma - note that tails may be a. Oligochaeta and Dipte	c groups: iy be damaged during sampling damaged during sampling ra)		Abundance 1-5 6-20 21-50 51-100 101+
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 =	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL-D (Gastropode <i>Asellus</i>	o the following 5 specifi ails) – note that tails ma - note that tails may be a. Oligochaeta and Dipte	c groups: iy be damaged during sampling damaged during sampling		Abundance 1-5 6-20 21-50 51-100 101+
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL-D (Gastropock A <i>sellus</i> ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroin/ertebrate gro		Abundance 1-5 6-20 21-50 51-100 101+
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 =	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL-D (Gastropock A <i>sellus</i> ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be s. Oligochaeta and Dipte aka and relative abunda <i>Ecdyonyrus</i> Ab	c groups: iy be damaged during sampling damaged during sampling ra)		Abundance 1-5 6-20 21-50 51-100 101+
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL-D (Gastropock A <i>sellus</i> ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be s. Oligochaeta and Dipte aka and relative abunda <u>Ecdyonurus</u> Ab <u>Rhithrogena</u> Ab	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroin/ertebrate gro	oup belo∴ (Abundance – Ap)	Abundance 1-5 6-20 21-50 51-100 101+ <i>Leuctra</i> Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL-D (Gastropock A <i>sellus</i> ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be - Oligochaeta and Dipte aka and relative abunda - Ecdyonurus Ab - Rhithrogena Ab - Haptagenia Ab	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroin/ertebrate gro	oup beloar (Abundance Ap) Pro	Abundance 1-5 6-20 21-50 51-100 101 + Leactra Ab Isoper/a Ab tonemura Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL-D (Gastropock A <i>sellus</i> ne total number of t	o the following 5 specifi ails) – note that tails may - note that tails may be s. Oligochaeta and Dipte aka and relative abunda <i>Ecdyonyrus</i> Ab <i>Rhithrogena</i> Ab <i>Haptagenia</i> Ab <i>Ephemerella</i> Ab	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroin/ertebrate gro	oup beloar (Abundance Ap) Pro	Abundance 1-5 6-20 21-50 51-100 101 + Leuctra Ab Isoperla Ab tonemura Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode <i>Asellus</i> ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithrogena</i> Ab <i>Haptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroin/ertebrate gro	oup beloar (Abundance Ap) Pro	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tonemura Ab phinemura Ab Perla Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropock Asellus ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdy-onurus</i> Ab <i>Rhithrogena</i> Ab <i>Heptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroin/ertebrate gro	pup belo.: • (Abundance – Ap) Pro -ן זוגר	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab Isoperla Ab chnemura Ab Perla Ab Dinocras Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropock Asellus ne total number of t	o the following 5 specifi ails) – note that tails may be - note that tails may be - Oligochaeta and Dipte aka and relative abunda - Ecdyonyrus Ab - Rhithrogena Ab - Heptagenia Ab - Caenis Ab - Caenis Ab - Caenis Ab - Caenis Ab	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroin/ertebrate gro	oup beloar (Abundance — An) <i>Pro</i> 4 <i>mp</i> Oth	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tonemura Ab tonemura Ab Perla Ab Perla Ab Denocras Ab her Piecop Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropock Asellus ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdy-onurus</i> Ab <i>Rhithrogena</i> Ab <i>Heptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab	c groups: y be damaged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera:	pup belo (Abundance – Ap) Pro վոր Oth Oth	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tonemura Ab whinemura Ab Perla Ab Dinocras Ab er Piecop Ab er Piecop Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode <i>Asellus</i> ne total number of t	o the following 5 specifi ails) – note that tails may be - note that tails may be - Oligochaeta and Dipte aka and relative abunda - Ecdyonyrus Ab - Rhithrogena Ab - Heptagenia Ab - Caenis Ab - Caenis Ab - Caenis Ab - Caenis Ab	c groups: y be damaged during sampling damaged during sampling (a) nce of each macroinvertebrate gro Plecoptera:	oup beloar (Abundance – Ao) Pro 1712 Oth Oth Total Relative	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tonemura Ab whinemura Ab Perla Ab Dinocras Ab er Piecop Ab er Piecop Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 = Calculate th	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode <i>Asellus</i> ne total number of t	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithrogena</i> Ab <i>Haptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab Paraleptophlebia Ab Other Ephem Ab Cher Ephem Ab	c groups: be damaged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Total no, of Taxa Lymnaea (G) Ab	oup beloar (Abundance – Ap) Pro 4712 Oth Oth Total Relative Chironomidae (D) Ab	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tonemura Ab whinemura Ab Perla Ab Dinocras Ab er Piecop Ab er Piecop Ab
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode Asellus ne total number of t <i>E</i> Xa O Total R	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte axa and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithrogena</i> Ab <i>Heptagenia</i> Ab <i>Caenis</i> Ab <i>Caenis</i> Ab <i>Caenis</i> Ab Other Ephem Ab Cother Ephem Ab Elative Abundance dae Ab G.OL.D:	c groups: by be damaged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Total no. of Taxa Ly mnaea (G) Ab Potamopy rgus (G) Ab	Dup beloar (Abundance Ab) Pro Imp Oth Oth Total Relative <u>Chironomidae (D) Ab</u> <u>Chironomidae (D) Ab</u>	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tonemura Ab binnemura Ab Perla Ab Perla Ab Perla Ab Perlecop Ab er Piecop Ab Abundance Asellus: Absent
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2 tails) Trichoptera G OL.D (Gastropock <i>Asellus</i> ne total number of t <i>Ka</i> <i>Materia</i> <i>Ka</i> <i>Materia</i> <i>Ka</i> <i>Materia</i> <i>Ka</i> <i>Materia</i> <i>Ka</i> <i>Materia</i> <i>Ka</i>	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithrogena</i> Ab <i>Haptagenia</i> Ab <i>Caenis</i> Ab <i>Caenis</i> Ab <i>Caenis</i> Ab Other Ephem Ab Other Ephem Ab telative Abundance dae Ab GOLD:	c groups: by be damaged during sampling damaged during sampling ra) nce or each macroin/entebrate gro Plecoptera: Total no. of Taxa Ly mnaea (G) Ab Potamopy rgus (G) Ab Planorbis (G) Ab	oup belo: (Abundance — Ap) Pro 1mp Oth Oth Oth Chironomidae (D) Ab Chironomus (D) Ab Simuliidae (D) Ab 3	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tomentura Ab Perla Ab Perla Ab Perla Ab Perla Ab Perlecop Ab ar Piecop Ab Abundance Absent Few/Low
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G-OL.D (Gastropock <i>Asellus</i> ne total number of t E xa Total R Hydropsychi Polycentropodi <i>Rhyacop</i> Philopotam	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithroğena</i> Ab <i>Haptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab Paraleptophlebia Ab Other Ephem Ab Caenis Ab Other Ephem Ab Etelative Abundance dae Ab <i>GoOL.D</i> : dae Ab	c groups: ay be damaged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Total no. of Taxa Ly mnaea (G) Ab Potamopy rgus (G) Ab Planorbis (G) Ab Ancy-lus (G) Ab	Dup beloa (Abundance – Ap) Pro 1mp Oth Oth Chironomidae (D) Ab Chironomus (D) Ab Simuludae (D) Ab Dicranota (D) Ab	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tomentura Ab Perla Ab Perla Ab Perla Ab Perlecop Ab ar Piecop Ab Abundance Absent Few/Low Common/
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropock <i>Asellus</i> ne total number of t E xa Total R Hydropsychi Polycentropodi <i>Rhyacop</i>	o the following 5 specifi ails) – note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithroğena</i> Ab <i>Haptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab Paraleptophlebia Ab Other Ephem Ab Caenis Ab Other Ephem Ab Etelative Abundance dae Ab <i>GoOL.D</i> : dae Ab	c groups: ay be damaged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Total no. of Taxa Lymnaea (G) Ab Potamopy rgus (G) Ab Planorbis (G) Ab Ancy-lus (G) Ab Physa (G) Ab	pup beloar (Abundance – Ap) Pro Imp Oth Oth Chironomidae (D) Ab Chironomidae (D) Ab Simuliidae (D) Ab Dicranota (D) Ab Tipulidae (D) Ab	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tomentura Ab Perla Ab Perla Ab Perla Ab Perla Ab Perlecop Ab ar Piecop Ab Abundance Absent Few/Low
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode <i>Asellus</i> ne total number of t E xa Total R Hydro <u>p</u> sychi Polycentropode <i>Rhyacop</i> Philopotam Limnephil Sericostomat	o the following 5 specifi ails) - note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithrogena</i> Ab <i>Haptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab Other Ephem Ab Other Ephem Ab Other Ephem Ab Caenis Ab Other Ephem Ab Caenis Ab Other Abundance dae Ab Chila Ab Caenis Ab	c groups: ay be damaged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Plecoptera: Total no. of Taxa Ly mnaea (G) Ab Potamopy rgus (G) Ab Planorbis (G) Ab Physa (G) Ab Lumbriculus (OI) Ab 5	Dup belos: (Abundance – Ap) Pro Imp Othe Othe Chironomidae (D) Ab Chironomidae (D) Ab Simuliidae (D) Ab Dicranota (D) Ab Tipulidae (D) Ab Chironomidae (D) Ab	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tonemura Ab Perla Ab Perla Ab Perla Ab Dinocras Ab er Piecop Ab Abundance Absent Few/Low Common, Numerous
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode <i>Asellus</i> ne total number of t E xa Total R Hydro <u>p</u> sychi Polycentropode <i>Rhyacop</i> Philopotam Limnephil Sericostomat Glossosomat	o the following 5 specifi ails) - note that tails ma - note that tails may be a Oligochaeta and Dipte aka and relative abunda <i>Ecdyonurus</i> Ab <i>Rhithrogena</i> Ab <i>Haptagenia</i> Ab <i>Ephemerella</i> Ab <i>Caenis</i> Ab Other Ephem Ab Other Ephem Ab Cher Ephem Ab	c groups: an aged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Plecoptera: Containo, of Taxa Lymnaea (G) Ab Potamopy rgus (G) Ab Planorbis (G) Ab Ancy-lus (G) Ab Physa (G) Ab Lymbriculus (OI) Ab Eiseniella (OI) Ab	pup beloar (Abundance – Ap) Pro Imp Oth Oth Chironomidae (D) Ab Chironomidae (D) Ab Simuliidae (D) Ab Dicranota (D) Ab Tipulidae (D) Ab	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tomentura Ab Perla Ab Perla Ab Perla Ab Perlecop Ab ar Piecop Ab Abundance Absent Few/Low Common/
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode <i>Asellus</i> ne total number of t E xa O Total R Hydropsychi Polycentropode <i>Rhy acop</i> Philopotam Limnephil Sericostomat Giossosomat Lepidostomat	o the following 5 specificalls) - note that tails may be a cligochaeta and Dipte as a and relative abunda <i>Ecdyonurus</i> Ab <i>Ecdyonurus</i> Ab <i>Haptagenia</i> Ab <i>Caenis</i> Ab <i>Caeni</i>	c groups: ay be damaged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Plecoptera: Total no. of Taxa Ly mnaea (G) Ab Potamopy rgus (G) Ab Planorbis (G) Ab Physa (G) Ab Lumbriculus (OI) Ab 5	Dup belos: (Abundance – Ap) Pro Imp Othe Othe Chironomidae (D) Ab Chironomidae (D) Ab Simuliidae (D) Ab Dicranota (D) Ab Tipulidae (D) Ab Chironomidae (D) Ab	Abundance 1-5 6-20 21-50 51-100 101+ <i>Leuctra</i> Ab <i>Isoperla</i> Ab <i>tonemura</i> Ab <i>conemura</i> Ab <i>Perla</i> Ab <i>Perla</i> Ab <i>Dinocras</i> Ab er Piecop Ab Abundance Asellus: Absent Few/Low Common, ¹ Numerous
The macroinverteb Group 1 = Group 2 = Group 3 Group 4 Group 5 Calculate th Ephemeroptera:	rates are divided int Ephemeroptera (3-t Plecoptera (2-tails) Trichoptera G OL.D (Gastropode <i>Asellus</i> ne total number of t E xa Total R Hydropsychi Polycentropodi <i>Rhyacop</i> Philopotam Limnephil Sericostomat Giossosomat Lepidostomat Other Trichopt	o the following 5 specificalls) - note that tails may be a cligochaeta and Dipte as a and relative abunda <i>Ecdyonurus</i> Ab <i>Ecdyonurus</i> Ab <i>Haptagenia</i> Ab <i>Caenis</i> Ab <i>Caeni</i>	c groups: an aged during sampling damaged during sampling ra) nce of each macroinvertebrate gro Plecoptera: Plecoptera: Containo, of Taxa Lymnaea (G) Ab Potamopy rgus (G) Ab Planorbis (G) Ab Ancy-lus (G) Ab Physa (G) Ab Lymbriculus (OI) Ab Eiseniella (OI) Ab	Dup belos: (Abundance – Ap) Pro Imp Othe Othe Chironomidae (D) Ab Chironomidae (D) Ab Simuliidae (D) Ab Dicranota (D) Ab Tipulidae (D) Ab Chironomidae (D) Ab	Abundance 1-5 6-20 21-50 51-100 101+ Leuctra Ab Isoperla Ab tornemura Ab Perla Ab Perla Ab Perla Ab Perla Ab Perlecop Ab Abundance Absent Few/Low Common, Numerous NOTE: Asee must be

e u tt

at TŐ

1

Step 1. Calculate the Index Score by circling the appropriate box representing the total number of taxa and the total abundance calculated from *each macroinvertebrate group* calculated from page 1 of the recording sheet and enter in to the boxes in Step 2.



Step 3. Calculate the Total Index Score, the Average Index Score and the SSR Score using the boxes below



SSR Score (AIS:x 2) O·8

Step 4. Assess the stream by comparing the final SSR score with the categories below and tick the appropriate box

04

> 7.25 Prebably not at risk	> 6.5 - 7.25 Indeterminate Stream may bellatirisk	< 6.5 Stream at risk	
Surveyor (signed):	Name (print):	Date:	1